



## Quantum Mechanics 3 2001/2002

### Solution set 2

(1)

(a) Stationary state means that the wavefunction factorizes:  $\psi(x, t) = u(x) \exp(-iEt/\hbar)$ . This is the simple time dependence for a state of definite energy. Whenever you see just  $u(x)$ , think of this as  $\psi(x, t = 0)$  – so that there will always be an implicit factor  $\exp(-iEt/\hbar)$  for the time dependence.

The full Schrödinger equation is  $-(\hbar^2/2m)\nabla^2\psi + V\psi = i\hbar\partial\psi/\partial t$ . Putting in the stationary state expression converts the rhs to just  $E\psi$ , giving the time-independent Schrödinger equation. Notice that the full Schrödinger equation allows superposition of solutions, but the time-independent form does not, because each state has a different energy: each state solves a different time-independent equation.

(b) Therefore, when you see a superposition like  $[u_1(x) + u_2(x)]/\sqrt{2}$ , this must be the  $t = 0$  form of the time-dependent superposition  $[\psi_1(x, t) + \psi_2(x, t)]/\sqrt{2}$ . At a later time, the full expression for  $\psi$  is

$$\psi = [u_1(x) \exp(-iE_1t/\hbar) + u_2(x) \exp(-iE_2t/\hbar)]/\sqrt{2},$$

and it is crucial that the two energies are different. The complex conjugate is

$$\psi^* = [u_1^*(x) \exp(+iE_1t/\hbar) + u_2^*(x) \exp(+iE_2t/\hbar)]/\sqrt{2},$$

so

$$|\psi|^2 = |u_1|^2/2 + |u_2|^2/2 + (u_1^*u_2/2) \exp[i(E_1 - E_2)t/\hbar] + (u_1u_2^*/2) \exp[-i(E_1 - E_2)t/\hbar].$$

If  $u_1^*u_2 = |u_1u_2| \exp(i\phi)$ , then

$$|\psi|^2 = |u_1|^2/2 + |u_2|^2/2 + |u_1u_2| \cos[(E_1 - E_2)t/\hbar + \phi].$$

In other words, the probability density oscillates back and forth. Quantum interference changes the probability density as a function of time.

(2)

(a) The Schrödinger equation for this case is

$$\left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + m\omega^2 x^2/2\right) \psi - E\psi = 0$$

(b) If  $\psi = Ae^{-\alpha x^2}$ , then the first 2 derivatives are

$$\begin{aligned}\psi' &= -2\alpha x Ae^{-\alpha x^2} \\ \psi'' &= -2\alpha e^{-\alpha x^2} + 4\alpha^2 x^2 Ae^{-\alpha x^2}.\end{aligned}$$

The resulting Schrödinger equation must be true for all  $x$ , so the total  $x^2$  coefficient in the equation must be zero:  $-2\hbar^2\alpha^2/m + m\omega^2/2 = 0$ , so  $\alpha = m\omega/2\hbar$ . Similarly, the constant term in the equation must vanish:  $\hbar^2\alpha/m - E = 0$ , so  $E = \hbar\omega/2$  - which proves the  $n = 0$  case of  $E = (n + 1/2)\hbar\omega$ .

(3)

(a) Parity:  $\psi(-x) = \pm\psi(x)$  (can always assume definite parity for a symmetric potential). Between the delta-potentials,  $V = 0$ , so the allowed wavefunctions are  $\psi \propto \exp(\pm\beta x)$ , where  $\beta^2 = -2mE/\hbar^2$ . here, we want a bound state, so  $E$  is negative and  $\beta$  is a real number. The positive parity solution therefore is

$$\psi = \begin{cases} Ae^{\beta x} & x < -a \\ B(e^{\beta x} + e^{-\beta x}) & -a < x < a \\ Ae^{-\beta x} & x > a \end{cases}$$

and for negative parity

$$\psi = \begin{cases} Ae^{\beta x} & x < -a \\ B(e^{\beta x} - e^{-\beta x}) & -a < x < a \\ -Ae^{-\beta x} & x > a \end{cases}$$

(using the only 2 allowed waves to make combinations with the right symmetry). The waves between the spikes are obviously cosh or sinh functions.

(b) As in the notes, integrate the Schrödinger equation across a delta function of weight  $A$  to get  $\Delta\psi' = (2mA/\hbar^2)\psi$ . If you're worried about internal structure in the well, remember  $k$  inside is  $\propto \sqrt{V}$  when  $V$  is large. The phase change across the well is  $KL$ , which is  $\propto \sqrt{VL}\sqrt{L}$ . If we keep  $VL$  fixed and let  $L \rightarrow 0$ , this tend to zero. We can treat  $\psi$  as constant inside the well.

(c) The boundary condition in (b) gives the same info at  $x = \pm a$  by symmetry. Matching  $\psi$  and writing down the discontinuity in  $\psi'$  gives 2 equations which can be divided to eliminate  $A$  and  $B$ . We get

$$\frac{2m|\alpha|}{\hbar^2} - \beta = \begin{cases} \beta \tanh \beta a & \text{(even parity)} \\ \beta / \tanh \beta a & \text{(odd parity)} \end{cases}$$

Sketch the lhs and rhs of these expressions against  $\beta$ . There is always an intersection for the even case, but not for the odd case if  $1/a > 2m|\alpha|/\hbar^2$ .