

Physical characterization of the binary Edgeworth-Kuiper Belt object 2001 QT_297

David J. Osip (*Observatories of the Carnegie Institution of Washington, Las Campanas Observatory*)

S. D. Kern* and **J. L. Elliot*†** (*Massachusetts Institute of Technology, *Department of Earth, Atmospheric and Planetary Sciences, † Department of Physics*)

Abstract

Following our discovery of 2001 QT_297 as the second known binary Edgeworth-Kuiper Belt Object in October of 2001 [IAUC 7733], we have carried out additional high spatial resolution ground based imaging in October and November of 2001 and July, August, and September of 2002. Using the Raymond and Beverly Sackler Magellan Instant Camera (MagIC) on the Baade and Clay 6.5m telescopes at Las Campanas Observatory in Chile, we have obtained accurate astrometric and photometric measurements in the Sloan r', i', and g' filters. Superb seeing conditions and PSF fitting allow an accurate determination of the binary component separation and position angle over time as well as a detailed study of color and temporal variability of the individual components. Here we present a physical characterization of the individual components of 2001 QT_297 based on these astrometric, color and variability measurements. We find the primary to exhibit colors about 0.3 magnitudes redder than solar with no evidence for variability. The secondary component, however, exhibits strong variability (~0.6 magnitudes) with a best fit period of 4.7526 ± 0.0007 hr for a single peak lightcurve or 9.505 ± 0.001 hr for a dual peaked lightcurve. The colors measured for the secondary also suggest variability. Based on a preliminary orbit fit for the pair using observations spanning a one year arc, we are able to estimate a system mass of $\sim 3.2 \pm 1 \times 10^{18}$ kg and provide constraints to the surface albedo of 9-14% for assumed densities between 1 and 2 g/cm³

1. Introduction

Pluto was discovered in 1930, and rotational variability was well observed for several decades (cf. Stern 1992). However, it wasn't until 1978 that the ninth planet in the solar system was found to have a companion, Charon (Christy and Harrington, 1978). Subsequent study of this binary system, especially during a 5-year period of mutual eclipses and occultations, provided a detailed characterization of Pluto and Charon, including accurate estimates of their diameters, albedos, masses, and densities (Tholen and Buie, 1997; Young *et al.*, 2000). For solar-system bodies without a binary companion or natural satellite, accurate estimates of most of these physical characteristics require a close spacecraft encounter.

Until recently Pluto-Charon remained the only known binary system in the trans-Neptunian region. Pluto and Charon are large and close (currently < 32 AU) objects in comparison to the bodies thus far detected in the Edgeworth-Kuiper Belt (EKB). The Pluto-Charon system is thought to have formed via a catastrophic collision, reaccretion of the debris cloud, and relatively rapid tidal evolution to the presently observed locked state (where the orbital period is identical to Pluto's rotation period such that Pluto always shows the same face to Charon). A similarly evolved binary EKO with much smaller diameters (~ 100 - 300 km) would have tidally evolved in a short time period (Toth, 1999; Weidenschilling *et al.*, 1989) to a separation of approximately 2000 km, which would subtend an angle of only 0.06 arcsec at a distance of 44 AU. Such a pair could not be resolved from any ground-based observatory (current adaptive optics systems are strictly only limited by the faint magnitude) and would provide a substantial challenge even utilizing the highest resolution available from the Hubble Space Telescope.

The previous argument would seem to limit our present study of the EKB to magnitudes, colors, and orbital elements. However, the first binary EKO, 1998 WW_31, was serendipitously discovered as a widely separated pair (angular separation of 1.2 arcsec) in re-analysis of images by C. Veillet and collaborators taken in December 2000 at the Canada France Hawaii Telescope (Veillet, 2001). In an October 2001 observing run that included additional physical studies of 1998 WW_31 and routine astrometry of EKOs discovered by the Deep Ecliptic Survey (Millis *et al.*, 2002), we made the unexpected discovery that 2001 QT_297 is also a binary (Elliot, 2001a,b). Our discovery observations (Figure 1) indicate a separation of only 0.61 arc seconds and a magnitude difference between the components ranging from approximately 0.6 to 0.9 magnitudes, implying a small ratio (~ 1.3) for the radii of the two bodies and providing initial evidence of variability.




Figure 1: Discovery observation of the binarity of 2001 QT_297. This single 300 second exposure was obtained with MagIC at the Magellan Project Baade 6.5m telescope on UT 11 October 2001 using a Sloan r' filter under exceptional seeing of 0.31 arc seconds.

2. Observations and Analysis

The discovery and all subsequent follow-up observations for this program were carried out with the Raymond and Beverley Sackler Magellan Instant Camera (MagIC) at the Magellan Project Baade and Clay 6.5m telescopes. MagIC is a highly efficient cryo-cooled CCD camera system utilizing a SITe 424a (2k x 2k) detector and equipped with a filter compliment including BVRI, and Sloan u',g',r',i',z'. Deployed at the f/11 focus of the Baade and Clay 6.5m telescopes, MagIC's field of view is 140 arc seconds on a side with a pixel scale of 0.069 arc seconds/pixel.

Following the discovery and confirmation observations on UT 11-12 October 2001, we have carried out 5 additional successful observing runs to investigate color and variability of the binary components and to establish an initial orbit fit to the high accuracy astrometry that we have obtained. Initial follow-up was carried out at the Baade telescope on UT 1-4 November 2001 using fractional nights during a previously scheduled instrument engineering run. The next successful observations were not until UT 13 July 2002, when time series photometry and color measurements were obtained (using the Sloan g',r',i' filters). An additional 7 hours of temporal monitoring was obtained (using Sloan r' and i' filters) on UT 18 July 2002. Following the successful July runs, astrometric observations were added on UT 7 August 2002 and 8 September 2002. Binary component separations and position angles are calculated from all the Sloan r' measurements for each night. These values are provided in Table 1 along with the measured time series differential photometry for the two binary components.

All frames were processed with standard overscan subtraction and flat fielding (using a combination of twilight and dome flats). For each night, the binary pair and a group of stellar comparisons were identified in each frame and standard circular aperture photometry was carried out to determine object centers, peak signal levels, and background levels. These parameters were then used as input for point spread function (PSF) modeling (Elliot *et al.* 1989; Bosh *et al.* 1993; Elliot *et al.* 2003). A roughly 2x2 arc second sub-array was selected from the calibrated image for each object, and an analytic PSF (in our case, either a circular Lorentzian or a circular

truncated Gaussian model (Schechter *et al.*, 1993) produced similar results) was fit to each object individually by least squares with the following free parameters: the object center (in row and column), the peak signal from the object, the background level, image diameter, and a parameter (“shape index”) that describes the shape of the PSF. Those objects that were confirmed point sources (similar diameter and shape index) were selected to form a set of comparison stars, and the brightest of these stars was defined to be the reference star for the field. Next a common PSF was fit to all of these stars, determining a common background, diameter and shape index. For the reference star, the free parameters were the peak signal, and position (row and column center). For each additional star in the fit, the ratio of its peak signal to that of the reference star plus row and column offsets from the reference star were fit. The binary components were fit as separate sources in this common fit with the initial position and peak value for the primary taken from the previous circular aperture photometry and for the secondary estimated from inspection of the calibrated frame. After the fitting was complete, the instrumental magnitude of the reference star, determined from the circular aperture photometry, and the fitted ratios for the other objects were used to calculate differential magnitudes for the target(s) and comparison stars.

Variability:

PSF profile fits to data on each of the nights are listed in Table 1. Clearly demonstrated in these data is a change in relative magnitude of the two components over the first two nights and to a lesser degree over the subsequent early follow-up observations in November. Careful comparison with multiple field stars on all images for Oct. 11 and 12 indicates the change in relative magnitude can be attributed entirely to variability of the secondary component. Such variability in solar system objects is generally regarded as evidence of rotation of either an elongated body or a surface with gross albedo differences (although interaction with a third body can not be ruled out). With evidence for variability of the secondary component, a concerted effort was made to carry out temporal monitoring of the pair during their next opposition in July 2002. The differential photometry results obtained on UT 18 July 2002 for the primary and secondary components of 2001 QT_297 and a check star are plotted in Figure 2. The primary component demonstrates little variability suggesting no rotation or a more spherical shape with no large-scale albedo variation. The secondary, however, shows strong variability of more than half a magnitude over the 7 hour observing interval.




Figure 2. Differential photometry plot from the night of UT 2002 July 18. The upper panel shows the magnitude difference between a comparison star and the 2001 QT_297 primary, the center panel plots the difference between the star and the secondary component, and the lower panel plots the difference between the comparison star and a check star.

The variability data for the UT 18 July 2002 observations suggests a possible single peak lightcurve of about 0.6 magnitude (peak to peak) amplitude with a period between 4.5 and 5 hours or a dual peaked lightcurve with twice that period. Since all evidence for variability can be attributed directly to the secondary component, we are able to use our complete data set of differential magnitude observations between the primary and secondary to determine a precise rotational period for this body. The advantage here is that the long time interval from October 2001 to July 2002 gives excellent leverage to the lightcurve phasing (i.e. with over 1200 rotations over the interval, a 3 second change in period will result in a one hour shift in phase). By first exploring phase dispersion minimization analysis of the entire data set and subsets on UT 13 and 18 July 2002, we were able to reduce the number of possible periodicities to explore. Additional explicit phasing of the entire data set was used to explore consistency between nights with recognizable slopes and turn-around points and to discriminate periodicities with contradictory data.




Figure 3. upper: Lightcurve phased to 4.7526 hr & lower: Lightcurve phased to 9.5055 hr. The most complete temporal sampling was achieved on UT 2002 July 18 (triangles), UT 2002 July 13 (diamonds) and UT 2001 Oct 11 (Xs), with other nights plotted using the (-) symbol. Data points and their associated uncertainties are plotted for all measurements in the Sloan r' filter tabulated in Table 1.

Our best phasing of the existing variability data, including a light time correction, leads to two preferred interpretations that are consistent with the current complete data set. Either the secondary has a rotation period of 4.7526 ± 0.0007 hr with a single peaked lightcurve (Figure 3 upper) presumably due to surface albedo/compositional variation; or the rotation period is 9.5055 ± 0.0010 hr for a dual peaked lightcurve (Figure 3 lower) indicating an elongated body with a semi-major to semi-minor axis ratio of approximately 1.8.

In either case, the implication remains that this was not a tidally evolved pair that underwent some gentle perturbation of the orbit to a more distant semi-major axis (i.e. via the effect of solar tides). Rather, the evidence for rapid variations is suggestive of the secondary component having undergone a subsequent impact that imparted both a change in orbital angular momentum to the system as well as rotational angular momentum to the body. Unfortunately the timescale for tidal evolution of the two components in their current orbit is longer than the age of the solar system and therefore not constraining on the time that such an impact occurred.

Colors:

Two color observations (using Sloan g' , r' , i' filters) were obtained for both binary components at a single phase on UT 13 July 2002. Applying zero-point magnitude and extinction corrections, calculated from a set of Sloan standard star observations, to the PSF fits for this night yields primary and secondary magnitude estimates listed in Table 2. While we make use of the Sloan filters for their well defined bands and high throughput, we have adopted the transformation equations of Fukugita *et al.*, 1996 to estimate the more often used BVRI colors for each component of the binary system. The primary component is consistently about three tenths of a magnitude redder than solar colors and well within the 'normal' range of small outer solar system bodies. At this phase, the secondary component colors are identical to the primary within the uncertainties.

Table 2. 2001 QT_297 colors

Object	Julian Date	g'	r'	i'	B-V ^a	V-R ^a	R-I ^a
Primary	2452468.785	23.79 ± 0.03	22.94 ± 0.02	22.55 ± 0.01	1.01 ± 0.04	0.61 ± 0.04	0.64 ± 0.03
Secondary	2452468.785	24.41 ± 0.07	23.55 ± 0.05	23.26 ± 0.05	1.00 ± 0.08	0.61 ± 0.07	0.59 ± 0.06
Secondary		del mag(r')- del mag(i') ^b		$r'-i'$	R-I		
	2452468.785	-0.01 ± 0.05		0.29 ± 0.05	0.53 ± 0.06		
	2452473.740	-0.02 ± 0.13		0.28 ± 0.13	0.52 ± 0.13		
	2452473.800	0.24 ± 0.10		0.54 ± 0.10	0.79 ± 0.10		
	2452473.868	0.13 ± 0.19		0.43 ± 0.19	0.68 ± 0.19		
	2452214.525	-0.09 ± 0.18		0.21 ± 0.18	0.45 ± 0.18		
	2452215.520	-0.01 ± 0.12		0.29 ± 0.12	0.53 ± 0.12		
	2452216.510	-0.41 ± 0.16		-0.11 ± 0.16	0.12 ± 0.16		

^a using transformations of Fukugita *et al.*, 1996 ^b differential magnitudes between primary and secondary

While accurate two color observations have only been obtained on one night thus far in our runs, we have obtained single color (Sloan r' and i') observations on five nights including very modest temporal coverage on UT 18 July 2002. Recognizing the variability of the secondary outlined in the preceding section and using the differential magnitudes for both the Sloan r' and i' filters tabulated in Table 1, our admittedly sparse data set is suggestive of possible strong color variation across the surface of the secondary component. Moreover, R-I colors (derived from the $r'-i'$ measurements) cover a magnitude range from 0.79 ± 0.10 to 0.12 ± 0.16 (Table 2 lower) indicating one of the reddest and the bluest EKO surface measured in these passbands (cf. Jewitt and Luu, 2001). Interestingly, the bluest measurements (recorded on UT 03 November 2001) correspond to at or near the peak brightness of the 2001 QT_297 secondary while the reddest measurements (recorded UT 18 July 2002) correspond to at or just preceding the minimum brightness of the secondary. This again hints toward an impact origin for the present orbit of this binary where the blue/bright region can be interpreted as fresher excavated material, the

reddest/faint region can be interpreted as a heavily irradiated unaltered surface with the remainder of the red surface being some combination of these two extremes.

3. Orbit

A preliminary orbit fit to the astrometric separations and position angles tabulated in Table 1 is diagrammed in Figure 4. The orbital parameters and their associated uncertainties are as follows: period = 876 ± 227 days, semi-major axis = $31,409 \pm 2500$ km, eccentricity = 0.31 ± 0.08 , inclination = 128.1 ± 6.5 degrees, node ascending longitude = 96.7 ± 13.4 degrees, argument of periapsis = 330.3 ± 22.4 degrees, and Epoch = 2001.787 ± 0.122 . The orbit fits are dependent on the final weighting of the measured positions and still suffer from an overall uniqueness problem due to the fact that the data acquired to date still consists of only two primary epochs (Oct-Nov 2001) and (Jul-Sep 2002). Additional observations will be needed to more tightly constrain the orbital parameters for this binary.




Figure 4. Orbit plot for the binary pair 2001 QT_297. The x-axis indicates the East-West positional offset in arc seconds of the secondary from the primary (at the origin). The y-axis indicates the North-South positional offset in arc seconds of the secondary from the primary. The points are a plot of the fit orbit and the crosses show the recorded positional offsets and their uncertainties.

Mass, Size, Albedo:

Using the current best estimates of orbital period and semi-major axis, we derived a system mass of $\sim 3.2 \times 10^{18}$ kg (while the uncertainties for the period and semi-major axis estimates are still large, they are correlated such that each feasible fit provides a similar system mass with an uncertainty of about $+0.3 \times 10^{18}$ and -0.2×10^{18} kg). Further adopting a mean magnitude difference between the two binary components of ~ 0.8 magnitudes and similar albedo and density values for the two bodies, we derived a mass ratio for the components of just over 3. Given the system mass and component mass ratio, we estimate size and albedo of the binary components under two plausible density conditions. First, assuming a density similar to that of Pluto (~ 2.0 g/cm³) we estimate the radius of the primary to be 78 km and that of the secondary to be 54 km with an albedo of 14%. At another extreme for a solid body, we assumed a density of 1.0 g/cm³ (for a composition of primarily water ice) and estimate primary and secondary radii of 98 and 68 km respectively with an albedo of 9%. Table 3 provides a comparison of the physical properties of 2001 QT_297 with those of the other EKO binary objects with known parameters, Pluto-Charon and 1998 WW_31. Of primary interest is the fact that the oft adopted EKO albedo estimate of 4% (based on measurements for a handful of cometary nuclei) does not appear a representative value for objects in the EKB unless in the unlikely case that their bulk densities are far lower than that of pure water ice. Additional evidence for the general inapplicability of cometary nuclei albedo values to EKO is found in combined visible and sub-mm observations of a handful of the largest EKO (eg. Jewitt *et al.*, 2001).

Table 3. EKO Binary Characteristics.

Binary	Period (days)	Semi-major axis (km)	Radius (km)	Albedo
Pluto-Charon ¹	6.38726 \pm 0.00007	19,636 \pm 8	1175 \pm 25 625 \pm 25	0.44-0.61 0.38
1998 WW_31 ²	574 +/- 10	22,300 +/- 800	118 & 98 ^a 148 & 123 ^b	0.09 0.07
2001 QT_297	876 +/- 227	31,409 +/- 2500	78 & 54 ^a 98 & 68 ^b	0.14 0.09

¹Tholen & Buie 1997 and references therein ²Veillet *et al.*, 2002 ^a density=2.0 g/cm³ ^b density=1.0 g/cm³

Continued resolved astrometric observations of 2001 QT_297 will allow us to refine the orbit parameters leading to a more accurate and precise mass determination for the system. Combined with accurate size and albedo estimates derived from simultaneous (or appropriately phased) visible observations of the individual components with thermal observations of the system with SIRTF, we will be able to uniquely determine the bulk density of this binary pair. Similar combinations of simultaneous visible and thermal observations of EKO should soon lead to accurate estimates of size and albedo for a statistically significant sample of these bodies as well as additional density estimates for the growing number of discovered binary systems.

This work was supported, in part, by NSF Grant AST99-77535 and NASA Grant NAG5-10444.

References

- Christy, J. W. and R. S. Harrington, 1978. The satellite of Pluto. *AJ* 83, 1005-1008
- Bosh, A. S., L. A. Young, J. L. Elliot, H. B. Hammel, and R. L. Baron 1992. Photometric variability of Charon at 2.2 μm . *Icarus* **95**, 319-324.
- Elliot, J. L., E. W. Dunham, R. L. Baron, A. W. Watts, S. P. Kruse, W. R. Rose, and C. M. Gillespie 1989. Image quality of the Kuiper Airborne Observatory. I. Results of the first flight series. *Publ. Astron. Soc. Pacific* **101**, 737-764.
- Elliot, J. L., D. J. Osip 2001, 2001 QT_297. IAUC 7765
- Elliot, J. L., S. D. Kern, D. J. Osip, S. M. Burles 2001, 2001 QT_297. IAUC 7733
- Elliot, J. L., M. Person, and S. Qu, 2003, Analysis of Stellar Occultation Data. II. Inversion, with application to Pluto and Triton. *AJ* 126 in press.
- Fukugita, M., T. Ichikawa, J. E. Gunn, M. Doi, K. Shimasaku, D. P. Schneider 1996, The Sloan Digital Sky Survey Photometric System. *AJ* 111, 1748-1756
- Jewitt, D. and J. Luu, 2001, Color and Spectra of Kuiper Belt Objects. *AJ* 122, 2099-2114
- Jewitt, D., H. Aousel, and A. Evans, 2001, The size and albedo of the Kuiper-belt object (20000) Varuna. *Nature* 411, 446-447
- Millis, R. L., M. W. Buie, L. H. Wasserman, J. L. Elliot, S. D. Kern, and R. M. Wagner 2002. The Deep Ecliptic Survey: A search for Kuiper Belt objects and Centaurs I. Description of methods and initial results. *Astron. J.* **123**, 2083-2109.
- Schechter, P. L., M. Mateo, and M. Saha 1993. DOPHOT, a CCD photometry program: Description and tests. *Publ. Astron. Soc. Pacific* **105**, 1342-1353.
- Tholen, D. J. and M. W. Buie, 1997, Pluto and Charon, 193-219
- Stern, S. A. 1992, The Pluto-Charon System. *Annu. Rev. Astron. Astrophys.* 30: 185-233.
- Toth, I., 1999, NOTE: On the Detectability of Satellites of Small Bodies Orbiting the Sun in the Inner Region of the Edgeworth-Kuiper Belt. *Icarus* 141, 420-425
- Veillet, C., A. Doressoundiram, J. Shapiro, J. J. Kavelaars, A. Morbidelli 2001, S/2000 (1998 WW_31) 1 IAUC 7610
- Veillet, C., J. Parker, I. Griffin, B. Marsden, A. Doressoundiram, M. Buie, D. Tholen, M. Connelley, M. Holman 2001, The binary Kuiper-belt object 1998 WW_31. *Nature* 416, 711-713
- Weidenschilling, S. J., P. Paolicchi, and V. Zappala, 1989, Asteroids II, 643-658
- Young, E. F., R. P. Binzel, and K. Crane, 2000, A Two-Color Map of Pluto's sub-Charon Hemisphere. *AJ* 121, 552-561

Table 1. 2001 QT_297 Photometry and Astrometry

Udate	JulianDate	delta_mag	Sloan_filter	Separation(")	position_angle°
2001-Oct-11				0.606 ± 0.005	117.2 ± 0.5
	2452193.50875	0.88 ± 0.04	r'		
	2452193.54084	0.64 ± 0.03	r'		
2001-Oct-12	2452193.56951	0.58 ± 0.02	r'		
				0.608 ± 0.008	116.1 ± 0.8
2001-Nov-01	2452194.57804	0.90 ± 0.04	r'		
				0.66 ± 0.02	108.9 ± 2.1
	2452214.501725	0.71 ± 0.17	r'		
	2452214.510515	0.73 ± 0.23	r'		
	2452214.519235	0.53 ± 0.14	r'		
	2452214.529020	0.62 ± 0.17	i'		
	2452214.536645	0.73 ± 0.20	i'		
	2452214.544280	0.54 ± 0.12	i'		
	2452214.552600	0.72 ± 0.12	i'		
2452214.562565	0.59 ± 0.18	i'			
2001-Nov-02				0.67 ± 0.02	105.9 ± 2.3
	2452215.497845	0.76 ± 0.19	r'		
	2452215.507160	0.36 ± 0.21	r'		
	2452215.515540	0.71 ± 0.12	r'		
	2452215.525000	0.72 ± 0.11	i'		
2001-Nov-03	2452215.544020	0.76 ± 0.10	r'		
				0.67 ± 0.04	106.7 ± 3.3
	2452216.506450	0.23 ± 0.14	r'		
2001-Nov-04	2452216.513250	0.64 ± 0.15	i'		
	2452216.520910	0.69 ± 0.08	i'		
				0.66 ± 0.02	102.1 ± 3.1
	2452217.495250	0.24 ± 0.24	r'		
2002-Jul-13	2452217.501065	0.45 ± 0.09	r'		
	2452217.572790	0.78 ± 0.26	r'		
	2452217.581665	0.81 ± 0.46	r'		
				0.76 ± 0.03	335.6 ± 1.6
	2452468.715270	1.00 ± 0.05	r'		
	2452468.762605	0.74 ± 0.08	r'		
	2452468.768470	0.80 ± 0.06	i'		
2452468.774115	0.67 ± 0.05	i'			
2452468.783717	0.62 ± 0.12	g'			
2452468.793260	0.66 ± 0.05	r'			
2452468.814015	0.54 ± 0.07	r'			
2452468.834775	0.64 ± 0.04	r'			
2002-Jul-18				0.78 ± 0.05	333.8 ± 5.3
	2452473.634430	0.99 ± 0.29	r'		
	2452473.668605	1.24 ± 0.15	r'		
	2452473.709265	0.84 ± 0.10	r'		

Draft

	2452473.736175	0.75 ± 0.11	r'		
	2452473.743860	0.77 ± 0.10	i'		
	2452473.771090	0.66 ± 0.15	r'		
	2452473.797980	0.73 ± 0.19	r'		
	2452473.805605	0.49 ± 0.09	i'		
	2452473.832620	1.00 ± 0.09	r'		
	2452473.864590	1.27 ± 0.19	r'		
	2452473.872205	1.14 ± 0.07	i'		
	2452473.897450	0.81 ± 0.10	r'		
	2452473.932750	0.70 ± 0.08	r'		
2002-Aug-07				0.92 ± 0.13	332.2 ± 8.4
	2452493.690120	0.87 ± 0.29	r'		
2002-Sep-08				0.93 ± 0.07	315.0 ± 6.6
	2452525.740135	0.94 ± 0.19	r'		