

Optimal capture of non-Gaussianity with weak lensing


Joel Bergé (JPL/Caltech)

Adam Amara (ETH Zurich)

Alexandre Réfrégier (CEA Saclay)

Why capture non-Gaussianity ?


- To break degeneracies left by the power spectrum



- How do we break those degeneracies ?

Weak lensing statistics

Jain, Seljak & White 1997 CDM model, $0.5 \times 0.5 \text{ deg}^2$



2-point correlation
function (power
spectrum) : $\langle \gamma_i \gamma_i \rangle$


3-point correlation
function (bispectrum):
 $\langle \gamma_i \gamma_j \gamma_k \rangle$

Galaxy clusters counts


Sensitive to non-
Gaussianity

Weak lensing as a cosmological probe : power spectrum and bispectrum

Power spectrum




Bispectrum




Weak lensing as a cosmological probe : galaxy clusters counts

$$\frac{d^2 N(z, w)}{d\Omega dz} = \frac{d^2 V_c}{d\Omega dz} \int_{M_{\text{lim}}}^{\infty} n(M, z, w) dM$$

$$\nu = \frac{\sqrt{n_g}}{\sigma_\gamma} \sqrt{\int d^2 x \kappa^2(x)}$$



Horellou & Bergé 2005



Capturing non-Gaussianity: high-precision cosmology

Bergé, Amara & Réfrégier 2010

- Fisher matrix analysis to forecast errors on cosmological parameters
- Halo model
- Euclid-like survey : 20,000 deg², 40 gal/arcmin², $z_m=1$
- Vary 8 parameters : Ω_m , Ω_Λ , Ω_b , σ_8 , h , n , w_0 , w_a
- Marginalized errors from :
 - weak lensing power spectrum
 - weak lensing bispectrum
 - weak-lensing-selected cluster counts
- We investigate cluster counts with different amount of information
 - S/N
 - redshift
 - mass
 - redshift and mass

Resolved vs unresolved parts


Discrimination against

- *resolved* part : $S/N > \nu_{th}$; non-linear part of data
- *unresolved* part : $S/N < \nu_{th}$; linear part of data

Massey et al 2007


Resolved part : cluster counts

Unresolved part : power spectrum




Resolved part

Unresolved part




Forecasted errors : Ω_m - σ_8 plane



Forecasted errors : dark energy plane

$$w(a) = w_0 + (1 - a)w_a$$




Forecasted errors : dark energy plane

Power spectrum

Power spectrum + Bispectrum

Power spectrum + Peak
counts (S/N>5)



Conclusions

- Cluster counts : counting S/N peaks breaks power spectrum degeneracies as well as counting clusters with additional information
- Bispectrum : provides constraints similar to those from the power spectrum
- Bispectrum : provides constraints similar to those from cluster counts (when combined with the power spectrum)

Guidelines to constrain cosmology with weak lensing

- 1) Measure power spectrum
- 2) Count S/N peaks, combine with 1)
- 3) Measure bispectrum, combine with 1), should be consistent with 2)