



Galaxies within the Epoch of
Reionization with
JWST/NIRSpec

Andrew Bunker



*On behalf of the
NIRSpec
Instrument Science
Team*




UNIVERSITY OF
OXFORD

NIRSpec


- Developed by the European Space Agency with Astrium GmbH and GSFC
- Operating wavelength: 0.6 – 5.0 μm
- Spectral resolution: 100, 1000, 3000
- Field of view: 3.4 x 3.4 arc minutes
- Aperture control: programmable micro-shutters, 250,000 pixels
- Angular resolution: shutter open area 203 x 463 mas, pitch 267 x 528 mas
- Detector type: HgCdTe, 2048 x 2048 pixel, 2 detectors, $T_{\text{op}} = 37\text{K}$ (passive)
- Reflective optics, SiC structure and optics



Multiple Objects
 ≤ 100 objects




NIRSpec microshutters



From STScI Newsletter, 2014, Karakla et al.

Spectra from NIRSpec



From STSci Newsletter, 2014, Karakla et al.

NIRSpec Instrument Science Team

Santiago
Arribas

Hans-Walter
Rix

Peter
Jakobsen


Harvey
Moseley

Bernie
Rauscher

Marijn
Franx

Roberto
Maiolino

...and Pierre Ferruit (ESA PI), Andy Bunker,
Stephane Charlot, Chris Willott



NIRSpec Galaxy Assembly: An expanding team...


Santiago Arribas, Andy Bunker, Bernd Huseman,
Stéphane Charlot, Chris Willott,
Emma Curtis-Lake, Marijn Franx,
Giovanna Giardino, Jacopo Chevallard,
Michael Maseda, Pierre Ferruit,
Peter Jakobsen, Roberto Maiolino,
Hans-Walter Rix, Stefano Carniani, Tim Rawle

NIRSpec GTO

MSA galaxy assembly program

Top priority science cases:


- Spectroscopic confirmation of the most distant galaxies
- Escape of Ly continuum and Ly α photons
- Stellar populations, kinematics and chemical enrichment
- PopIII
- AGNs and their host galaxies
- Large scale structure, environment
- Evolution of the joint distributions



Wilkins et al.
(2010) MNRAS
The Luminosity
Function at
 $z \sim 7$

An increasing problem for reionization: requires steep faint-end slope ($\alpha < -1.7$), large contribution from unobserved faint galaxies, high escape fraction ($f_{\text{esc}} > 0.2$) and very smooth IGM (low clumping, $C < 5$)

Received on earth, after
passing through IGM



"Lyman break technique" - sharp drop in flux at λ below
Ly- α . First done by Steidel et al. have >1000 $z \sim 3$
objects, "drop" in U-band. Now with HST/WFC3 in the
near-IR pushing "optical dropout" galaxies at $z \sim 6-12$. At
 $z > 6$ almost complete Gunn-Peterson absorption below
Lyman-alpha.

Overall NIRSpec Galaxy Assembly Plan

a wedding cake survey at $R=100$ and $R=1000$, about 500hours

- Deep, 20-40 sq arcmin, 1-5 μm , 40-45% of the time, $AB \leq 29-30$, $2 < z < 14$
- Medium, 100-200 sq arcmin, 1-5 μm , 40-45% of the time, $AB \leq 27-28$, $2 < z < 14$
- Shallow, >400 sq arcmin, 2-5 μm , 10-20% of the time,
 $AB \leq 25-26$, 7000+ spectra, $2 < z < 4$ ($4 < z < 14$)

Fields: deep supporting observations (e.g., MIPS, Herschel, ALMA, X-Ray, ...):

HUDF, HDF, GOODS-N,S, CANDELS for shallow, Frontier Fields, Clash

For Deep, probably HUDF and GOODS-North (in collaboration with NIRSpec)

4) $R=3000$ IFU spectroscopy of extended objects ≈ 300 hours

Deep

- Per field: integrations: $R=100$, $1e5$ seconds
 - $R=1000$, 3 settings, total = $1e5$ seconds
 - Total integration time = $2e5$ seconds, with overheads 84 hours/field
- 2-3 pointings following up HST & NIRCAM deep fields
collaboration between NIRCAM & NIRSspec ISTs

Confirmation Highest redshift sources (1)

- **Science Objectives**

- Get spectroscopic redshifts at $z > 7$
- Initial characterization of spectra

- **Methods**




- Measure redshift from emission line/continuum features
- Simple modelling of the spectra (potentially pop III, continuum UV slopes, etc)

NIRSpec will get very accurate redshifts, and hence determine accurate rest-frame properties;

measure emission lines ($H\alpha$, $H\beta$, [OIII], [OII]), to constrain: attenuation by dust, star formation rate; ionization state and metallicity of the interstellar gas; presence of an AGN; HeII-1640 for pop III?

The Star Formation History of the Universe

I-drops in the Chandra Deep
Field South with HST/ACS
Stanway, Bunker & McMahon
2003



Confirmation of highest redshift (2)

- Are the blue spectral slopes real?
- Is the suggested severe contamination of broad-band magnitudes (e.g. Spitzer/IRAC) by huge EW line emission real?
- **Link to the program at large**
 - part of Deep UDF, Deep, Medium, Shallow;
 - $R=100$ and $R=1000$ to get emission lines
- **Ancillary Data:** HST UDF, NIRCcam imaging
 - Target selection (based on SED (photoz), dropout, etc, magnitude, size)
 - Morphological information (sizes)
 - photometry for accurate SED




The Lyman continuum escape fraction (1)

Science goals

- Escape fraction of Lyman continuum as a function of galaxy properties (mass, SFR, z ,....)

Methods

- Balmer lines compared with UV continuum
- Balmer decrement and UV slope -> dust extinction + age
- HeII/HeI/H line ratio -> shape of ionizing continuum
- $R = 1000$ ($R=100$ for Balmer break)





JWST-NIRSpec – Escape Fraction and Reionization

Escape fraction (from Balmer lines compared with UV)
(dependence on mass, SFR; compare with Ly α flux and profile)


The comparison of the H-alpha or H-beta photons with the UV continuum should provide a good constraint - a simple photon budget governed by well known photoionization and recombination physics.

[Modulo the UV extrapolation beyond Lyman-alpha, which may be model dependent, but can be constrained by the HeI/H ratio, or even HeII/HeI/H.]


Also: searches for PopIII – HeII 1640Ang



Ly-alpha fraction (Stark et al. 2010)



Lyman alpha equivalent width distribution as a function of redshift for patchy or not reionization scenarios. Also, measuring Lyman alpha line shape evolution into the reionization epoch with systemic redshifts determined from nebular lines.









Stark et al. (2010) Caruana et al. (2012, 2014). Ly-alpha suppressed in mostly-neutral IGM




Quantities to be measured:

- Stellar mass
- Galaxy age and star formation history
- Star Formation Rate
- Stellar metallicity
- Gas metallicity
- Attenuation by dust
- Presence and power of AGN
- Black hole masses
- Presence of shocks
- Outflows
- Dynamics

NIRSpec MSA simulations; $T_{exp} = 20,000$ sec $z=4.5$ with R100

NIRSpec team (Franx, Bunker, Ferruit, Maiolino, Arribas, Charlot, Rix, Willot, Starburst Jakobsen) Post-starburst





Absorption lines at $z > 5$ – few examples of bright Lyman break e.g. $z=5.5$ galaxy Dow-Hygelund et al (2005), AB=23-24, VLT spectrum (22 hours). Also VUDS (Le Fevre et al. 2015). R=1000 mode with NIRSspec to probe $z > 6$ rest-UV absorption lines

Synergies with other JWST instruments

NIRCAM:

- imaging (and slitless spectroscopy), target selection
- morphological parameters
- correction for slit losses
- calibration of physical parameters inferred from broad band photometry (e.g. photo-z calibration)

MIRI:

- mid-IR diagnostics of star formation and AGN
- complementary tracers of dust

NIRISS: complementary survey

- larger statistics for line emitters
- diagnostics limited to lower redshifts (blue part of the spectrum)

Conclusions

- Have found star-forming galaxies at $z=6-10$ (Lyman breaks), and spectroscopic confirmation at $z\sim 6$; not much Ly-alpha emission beyond $z\sim 7$ (due to Gunn-Peterson absorption?)
- However, $z>7$ number counts from HST/WFC3 imply the newly-discovered galaxies would struggle to reionize the IGM
- Many of these have very blue rest-UV spectral slopes
- High escape fraction/Steep faint end slope/low metallicity/smooth IGM?
- JWST spectroscopy will get $H\alpha$, $H\beta$, [OIII], [OII] to high redshift, Getting REAL redshifts for luminosity functions
- will determine escape fractions, star formation rates, metallicities...

Program plan – Medium Survey (100-200 sq. arcmin)

- 9 NIRSPEC pointings x 2 fields (test cosmic variance)
- (9arcmin x 9arcmin per field; possible fields include
- GOODS/CANDELS, Frontier Fields –
- complementary data, X-ray, UV, sub-mm....)
- ~3000 galaxies at $2 < z < 6$ (150 galaxies per pointing)
- $AB < 27$
- R=100: 20ksec per pointing (continuum)
- R=1000: 10ksec per pointing in each of three bands
(emission lines)
- total of ~ 230 hours

- Shallow survey (200-400 sq. arcmin, 48 pointings)

- $AB < 25$, 7000+ galaxies, R=100 for 1000s
- R=1000 at 2-5microns for 1000s (2 settings)
- Total of ~50 hours